

Species

To Cite:

Ayas D, Erguden SA, Erguden D. The occurrence of *Zu cristatus* (Family: Trachipteridae), in the Northeastern Mediterranean, Türkiye. *Species* 2024; 25: e28s1677
doi: <https://doi.org/10.54905/disssi.v25i75.e28s1677>

Author Affiliation:

¹Mersin University, Fisheries Faculty, Mersin, Turkey

²Vocational School of Imamoglu, University of Cukurova, Imamoglu, Adana, Türkiye

³University of Iskenderun Technical, Faculty of Marine Science and Technology, Iskenderun, Hatay, Turkey

Corresponding Author

University of Iskenderun Technical, Faculty of Marine Science and Technology, Iskenderun, Hatay, Turkey

Email: deniz.erguden@iste.edu.tr; derguden@gmail.com

ORCID List

Deniz Ayas	0000-0001-6762-6284
Sibel Alagoz Erguden	0000-0003-4363-433X
Deniz Erguden	0000-0002-2597-2151

Peer-Review History

Received: 04 April 2024

Reviewed & Revised: 08/April/2024 to 10/June/2024

Accepted: 13 June 2024

Published: 18 June 2024

Peer-Review Model

External peer-review was done through double-blind method.

Species

pISSN 2319-5746; eISSN 2319-5754



© The Author(s) 2024. Open Access. This article is licensed under a [Creative Commons Attribution License 4.0 \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

The occurrence of *Zu cristatus* (Family: Trachipteridae), in the Northeastern Mediterranean, Türkiye

Deniz Ayas¹, Sibel Alagoz Erguden², Deniz Erguden^{3*}

ABSTRACT

On April 16, 2024, a scalloped ribbonfish *Zu cristatus* specimen with a standard length of 615 mm and a weight of 337.4 g was caught by a commercial trawler off Erdemli in the Gulf of Mersin (Northeastern Mediterranean, Türkiye) from 620-690 m depth range. The present paper reports the presence and is the first confirmation of *Z. cristatus* from the Gulf of Mersin, Türkiye (Northeastern Mediterranean). All measurements, counts, and color descriptions of *Z. cristatus* agree with previous descriptions. This record is a rare occurrence of the specimen of *Z. cristatus* on the Mediterranean coast of Türkiye. In addition, the detailed morphological measurements of *Z. cristatus* were given for the first time on the Turkish coast.

Keywords: Lampriformes, scalloped ribbonfish, deep-waters, Mersin Bay, Mediterranean Sea

1. INTRODUCTION

To date, two species of the family Trachipteridae have been reported from the Mediterranean Sea in shallow waters. These species are known as the Mediterranean dealfish *Trachipterus trachipterus* (Gmelin, 1789), and the scalloped ribbonfish *Zu cristatus* (Bonelli, 1819). The scalloped ribbonfish *Z. cristatus* belongs to the genus *Zu* and it is a meso-bathypelagic and cosmopolitan species distributed in the Mediterranean, Atlantic, Indian, and Pacific Oceans (Mundy, 2005). The species was first described in 1920 (Bonelli, 1920) as *Trachipterus cristatus*. The currently accepted and valid name of the species is *Z. cristatus*.

In the Adriatic and Mediterranean waters, *Z. cristatus* occurrence has been reported at different depths, from shallow waters in the Adriatic Sea Dulčić, (2002), Dulčić et al., (2014), Ionian Sea Mytilineou et al., (2013), Albano et al., (2022), Ligurian Sea Bonelli, (1820), Gavagnin, (1976), Fischer et al., (1987), Bottaro et al., (2005), Garibaldi, (2015), Tyrrhenian Sea Bianco et al., (2006), Psomadakis et al., (2006), Psomadakis et al., (2007), Zenetos et al., (2015), Falsone et al., (2017), Tiralongo et al., (2019), Tiralongo et al., (2020), Stipa et al., (2022), Sardina Cau, (1980), Balearic Sea García-Barcelona et al., (2014), the Iberian Sea Ibanez and Gallego, (1974), the western

Mediterranean between the coasts of Spain and Algeria Quignard and Tomasini, (2000), Catalonia Roig and Demestre, (1982) and in the Gulf of Tunis (Tunisia) Postel, (1955), Bradai and ElOuaer, (2012), the eastern Mediterranean in Israel Golani et al., (2023), Greece Papaconstantinou, (1998), Kaminas et al., (2021) and Turkey (Akyol and Tosunoğlu, 2023; Erguden et al., 2023).

In Turkish marine waters, *Z. cristatus* was first recorded in the Mediterranean Sea Akyüz and Artüz, (1957) and then in the Aegean Sea (Geldiay, 1969). Recently, this species has been reported from Saros Bay (northern Aegean Sea) by Akyol and Tosunoğlu, (2023) and from the Gulf of Antalya (Antalya and Finike coasts, eastern Mediterranean Sea) by (Çağıltay et al., 2023; Erguden et al., 2023). Although *Z. cristatus* Bilecenoğlu et al., (2002), Fricke et al., (2007) has been reported in Turkish checklists in Turkish marine waters, no record of this species has been reported from the northeastern Mediterranean coast of Turkey. Our present report is the first record of *Z. cristatus* in the Gulf of Mersin (Northeastern Mediterranean).

2. MATERIAL AND METHODS

A single specimen of *Z. cristatus* was caught by a commercial trawler on April 16, 2024, off Erdemli, Mersin Gulf in international waters (Coordinates: 36°04'202"N 34°31'597"E, 36°02'39.7"N 34°47'31.0" E) in the 620-690 m depth range (Figure 1). After the scalloped ribbonfish sample was taken from the commercial trawler where it was caught, it was taken to the Museum of Marine Life at Mersin University, where the species was identified and measured. The morphometric measurements of the specimen were made using a digital caliper and the weight was measured using a digital weight scale. All measurements and counts, as well as the morphological description and color, are consistent with the descriptions of (Bauchot, 1987; Olney, 1999). The specimen was registered at the Mersin University Marine Life Museum with catalog number MEUFC-24-11-148 and preserved in neutral formalin (Figure 2).




Figure 1 Map showing the fishing area of scalloped ribbonfish *Zu cristatus* in international waters off the Gulf of Mersin



Figure 2 The specimen of the scalloped ribbonfish *Zu cristatus* was captured (615 mm, SL) from the Gulf of Mersin, Türkiye

3. RESULTS

The main diagnostic characters and morphometric measurements of the captured specimen of *Z. cristatus* are given in centimeters: The specimen had a total length of 615 mm and a total weight of 337.4 g. The body is dorsoventrally compressed. The eye is large. The mouth is small and terminal, the first dorsal fin is highly elongated, the anal fin is absent, and paired pectoral and pelvic fins are present. In juveniles, the body is silvery, with approximately six vertical bars on the dorsal part of the body and four on the ventral part. Dark spots are on the body or fins. The dorsal fin has 120 rays and the pectoral fin has ten rays. The caudal fin rays could not be counted because our the tail sample was broken off was broken off during netting. Some morphometric measurements of the *Z. cristatus* specimen are given in (Table 1).

Table 1 Morphometric measurements of *Z. cristatus* in the North-Eastern Mediterranean

Metric	Value (mm)	% (SL)
Standard length	615	100
Head length	106	17.24
Pre-orbital length	36	5.85
Post-orbital length	41	6.67
Eye diameter	32	5.20
Operculum height	88	14.31
Upper jaw length	44	7.16
Lower jaw length	62	10.08
Pre-pectoral length	94	15.29
Pectoral fin length	43	6.99

Width of pectoral fin base	8	1.30
Maximum height of dorsal fin	48	7.81
Maximum height of the body	119	19.35
Total weight (g)	337.4	-
Meristic		
Lateral line scales	100	-
Pectoral fin rays	11	-
Dorsal fin spines	6	-
Upper jaw teeth	17	-
Lower jaw teeth	12	-
Palatine teeth	4	-
Vomerine teeth	3	-
Gill rakers total	10	-
Gill rakers epibranchial	2	-
Gill rakers ceratobranchial	8	-

4. DISCUSSION

In the Mediterranean, data on the biology and ecology of *Z. cristatus* are very limited. The scalloped ribbonfish *Z. cristatus* is an offshore fish with a maximum size of 118 cm SL (Heemstra and Kannemeyer, 1986). It feeds on pelagic crustaceans, small fish, and squid. Eggs are free-floating, large, and reddish. Juveniles typically swim at the surface, trailing elongated dorsal and pelvic fin rays that give them the appearance of a jellyfish. Juveniles specimens are characterized by a ribbon-shaped body, a short head, and a narrow mouth with a distinctly protruding upper jaw (Heemstra and Kannemeyer, 1986; Olney et al., 1993). *Z. cristatus* is distributed in tropical and temperate waters. This species is rarely recorded, especially in commercial trawls, and it is generally considered to be circumglobal in the epipelagic and mesopelagic zones from zero to 0-950 m depth (Fricke et al., 2011).

However, *Z. cristaus* is rarely found in the eastern Mediterranean. This species is mostly highly abundant in the central part of the Mediterranean Sea basin (Albano et al., 2022). In our study, the fish were caught in the deep waters of Mersin Gulf at approximately 690 m. Çağiltay et al., (2023) reported this species from shallower waters at 450 m in Antalya Bay. Akyol and Tosunoglu, (2023) reported an adult specimen of *Z. cristatus* caught at a depth of 600 m in Saros Bay (northern Aegean Sea, Turkey). Quigley and Henderson, (2014), reported that *Z. cristatus* that adults are commonly found at depths of 150-800 m, while juveniles prefer shallower waters. Golani et al., (2023) reported finding juveniles of this species near the surface in the eastern Mediterranean.

The young individual reported in this study was caught in deeper water than reported in the literature. In the present study, the occurrence a specimen of *Z. cristatus* was accidentally caught during commercial bottom trawling in Mersin Gulf. At the same time, this report represents the fourth occurrence of *Z. cristatus* on the Levantine Sea coast of Turkey. In addition, morphological measurements of *Z. cristatus* were reported for the first time in Turkish waters in this study. Although *Z. cristatus* has been classified as Least Concern (LC) on the IUCN Red List. This species remains Data Deficient (DD) in Turkey Fricke et al., (2007), this finding is significant as it contributes to our understanding of the distributional role of the species in the region.

5. CONCLUSION

We report the first substantiated record of *Z. cristatus* from the Gulf of Mersin (Northeastern Mediterranean coast, Türkiye). Although there is no evidence from past and present catch records that the species has established a permanent population along the northeastern Mediterranean coast, the fact that the individual reported in this study is a juvenile *Z. cristatus* provides important evidence for the permanent presence of the species in the northeastern Mediterranean. In addition, new records of the availability of scalloped ribbonfish along the Levantine coast indicate the successful existence of this species in recent years.

Acknowledgements

The authors thank the captain and boat staff for providing the fish sample.

Authors' Contribution

DE: Writing - original draft, methodology, writing - review and editing

SAE: Editing, final check control, validation

DA: Conceptualization, Investigation, sampling, visualization, original draft, writing - review and editing, validation

Ethical Approval

The Animal ethical guidelines are followed in the study for species observation & identification.

Informed consent

Not applicable.

Conflicts of interests:

The authors declare that there are no conflicts of interests.

Funding:

The study has not received any external funding.

Data and materials availability

All data associated with this study are present in the paper.

REFERENCES

1. Akyol O, Tosunoğlu Z. First occurrence of *Zu cristatus* (Trachipteridae) in the Turkish Aegean Sea (Eastern Mediterranean Sea). *Mar Life Sci* 2023; 5(1):37-39. doi: 10.5175 6/marlife.1285550
2. Akyüz E, Artüz I. Some observations on the biology of tuna (*Thunnus thynnus*) caught in Turkish waters. *GFCM Proc Tech Papers* 1957; 4:93-99.
3. Albano M, D'Iglio C, Spanò N, Di-Paola D, Alesci A, Savoca S, Capillo G. New report of *Zu cristatus* (Bonelli, 1819) in the Ionian Sea with an in-depth morphometrical comparison with all Mediterranean records. *Fishes* 2022; 7(6):305. doi: 10.3390/fishes7060305
4. Bauchot ML. Poissons osseux. (W. Fischer, M.L. Bauchot and M. Schneider, Eds.), Fiches FAO d'identification pour les besoins de la pêche. (rev. 1). Méditerranée et mer Noire. Zone de pêche 37. Commission des Communautés Européennes and FAO, Rome, 1987; 2:891-1421.
5. Bianco PG, Zupo V, Ketmaier V. Occurrence of the scalloped ribbonfish *Zu cristatus* (Lampridiformes) in coastal waters of the central Tyrrhenian Sea, Italy. *J Fish Biol* 2006; 68:150-155.
6. Bilecenoglu M, Taskavak E, Mater S, Kaya M. Checklist of the Marine Fishes of Turkey. *Zootaxa* 2002; 113:1-194. doi: 10.116 46/ZOOTAXA.113.1.1
7. Bonelli FA. Description d'une nouvelle espèce de poisson de la Méditerranée appartenant au genre Trachyptère avec des observations sur les caractères de ce même genre. *Mem Reale Accad Sci Torino* 1820; 24:485-494.
8. Bottaro M, Ferrando S, Ferrando T, Psomadakis PN, Vacchi M. Melanism in the gastric mucosa of the scalloped ribbonfish from the Ligurian Sea. *J Fish Biol* 2005; 66:1489-1492. doi: 10.11 11/j.0022-1112.2005.00680.x
9. Bradai MN, ElOuaer A. New record of the scalloped ribbonfish, *Zu cristatus* (Osteichthyes: Trachipteridae) in Tunisian waters (central Mediterranean). *Mar Biodiv Rec* 201 2; 5:e59. doi: 10.1017/S175526721200036X
10. Çağiltay F, Biçer E, Yıldız A, Gökoğlu M. Occurrence of the scalloped ribbonfish *Zu cristatus* (Lampridiformes) in the Gulf of Antalya, Türkiye. *NEsciences* 2023; 8(2): 82-85. doi: 10.2897 8/nesciences.1338121
11. Cau A. Second note on the bathyal ichthyofauna of the seas around southern Sardinia. *Quaderni della Civica Stazione Idrobiologia di Milano* 1980; 8:39-44.
12. Dulčić J, Dragičević B, Pavičić M, Ikica Z, Joksimović A, Markoić O. Additional records of non-indigenous, rare, and less known fishes in the eastern Adriatic. *Ann Ser Hist Nat* 2014; 24(1):17-22.
13. Dulčić J. First record of scalloped ribbon fish, *Zu cristatus* (Pisces: Trachipteridae), eggs in the Adriatic Sea. *J Plankton Res* 2002; 24(11):1245-1246. doi: 10.1093/plankt/24.11.1245
14. Erguden D, Erguden SA, Ayas D. A rare occurrence and confirmed record of scalloped ribbonfish *Zu cristatus*

- (Osteichthyes: Trachipteridae) in the Gulf of Antalya (Eastern Mediterranean), Turkey. *Ann Ser Hist Nat* 2023; 33(1):89-96. doi: 10.19233/ASHN.2023.13
15. Falsone F, Geraci ML, Scanella D, Okpala COR, Giusto GB, Bosch-Belmar M, Gancitano S, Bono G. Occurrence of two rare species from order. Lampriformes: Crestfish *Lophotus lacepede* (Giorna, 1809) and scalloped ribbonfish *Zu cristatus* (Bonelli, 1819) in the northern coast of Sicily, Italy. *Acta Adriat* 2017; 58(1):137-144. doi: 10.32582/aa.58.1.11
 16. Fischer W, Bauchot ML, Schneider M. Fiches FAO d'Identification des Espèces pour les Besoins de la Pêche. Méditerranée et Mer Noire, Volume II (Vertébrés). FAO, Rome, 1987; 1529.
 17. Fricke R, Bilecenoglu M, Sarı HM. Annotated checklist of fish and Lamprey species (Gnathostomata and Petromyzontomorphi) of Turkey, including a Red List of threatened and declining species. *Stuttgarter Beitr Naturk Sea A* 2007; 706:1-169.
 18. Fricke R, Kulbicki M, Wantiez L. Checklist of the fishes of New Caledonia, and their distribution in the Southwest Pacific Ocean (Pisces). *Stuttgarter Beitr Naturk A, Neu Serie* 2011; 4:341-463.
 19. García-Barcelona S, García-Cancela R, Cayuela MJ, Fernández-Peralta L, de Carlos A, Bañón R, Macías D, Báez JC. Descripción de dos ejemplares de *Zu cristatus* (Bonelli, 1820) capturados accidentalmente con un palangre semipelágico en el Mediterráneo occidental. *Arx Miscel Lania Zool* 2014; 14:91-98. doi: 10.32800/amz.2016.14.0091
 20. Garibaldi F. By-catch in the mesopelagic swordfish longline fishery in the Ligurian Sea (Western Mediterranean). *Collect Vol Sci Pap ICCAT* 2015; 71(3):1495-1498.
 21. Gavagnin EP. Considerazioni sulla cattura di uno *Zu cristatus* (Bonelli) a San Remo (Osteichthyes Trachipteridae). *Natura* 1976; 67:258-261.
 22. Geldiay R. Izmir Körfezinin başlıca balıkları ve muhtemel invasionları. Ege Üniversitesi Fen Fakültesi Monografileri, Izmir, 1969; 135 (in Turkish).
 23. Golani D, Edelist D, Morov AR, Stern N. The first confirmed record of *Zu cristatus* in the Mediterranean coast of Israel and the eastern shores of the Levant. *Mediterr Mar Sci* 2023; 24(1): 87-89. doi: 10.12681/mms.30976
 24. Heemstra PC, Kannemeyer SX. Trachipteridae. (M.M. Smith and P.C. Heemstra (Eds.), *Smiths' sea fishes*. Springer-Verlag, Berlin, 1986; 399-402.
 25. Ibanez M, Gallego L. A new record of a *Zu cristatus* (Trachipteridae, Pisces) off the coast of Blanes (Spain). *Vie et Milieu* 1974; 523-526.
 26. Kaminas A, Minasidis V, Doumpas, N, Naasan Aga Spyridopoulou RTF. Occurrence of *Trachipterus trachipterus* and *Zu cristatus* in the Greek Seas: Contribution of citizen science projects in rare species monitoring. In *Proceedings of the HydroMedit*, University of Thessaly, Thessaly, Greece, 2021.
 27. Mundy BC. Checklist of the fishes of the Hawaiian Archipelago. *Bishop Mus Bull Zool* 2005; 6:1-704.
 28. Mytilineou C, Anastasopoulou A, Christides G, Bekas P, Smith CJ, Papado-Poulou KN, Lefkaditou E, Kavadas S. New records of rare deep-water fish species in the Eastern Ionian Sea (Mediterranean Sea). *J Nat Hist* 2013; 47(25-28):1645-1662. doi: 10.1080/00222933.2013.775372
 29. Olney JE, Johnson GD, Baldwin CC. Phylogeny of lampridiform fishes. *Bull Mar Sci* 1993; 52:137-169.
 30. Olney JE. Order Lampriformes. (K.E. Carpenter and V.H. Niem, Eds.), *FAO Species Identification Guide for Fishery Purposes. The Living Marine Resources of the Western Central Pacific. Batoid Fishes, Chimaeras, and Bony Fishes Part 1 (Elopidae to Linophrynidae)*; FAO, Rome, Italy, 1999; 3: 952-959.
 31. Papaconstantinou C. Checklist of marine fishes of Greece, in *Fauna Graeciae*, 4, Athens, 1998; 257.
 32. Postel E. Capture d'un trachyptère *Trachipterus cristatus* Bonelli en Baie de Tunis. *Bull Inst Natl Sci Technol Mer* 1955; 51:69-71.
 33. Psomadakis PN, Scacco U, Vacchi M. Recent findings of some uncommon fishes from the central Tyrrhenian Sea. *Cybiuim* 2006; 30(4):297-304. doi: 10.26028/cybiuim/2006-304-002
 34. Psomadakis PN, Bottaro M, Vacchi M. On two large specimens of *Zu cristatus* (Trachipteridae) from the Gulf of Genoa (NW Mediterranean). *Cybiuim* 2007; 31(4):480-482. doi: 10.26028/cybiuim/2007-314-011
 35. Quigley DTG, Henderson G. The first record of the scalloped ribbonfish *Zu cristatus* (Bonelli, 1819) (Lampriformes: Trachipteridae) from N.W. European waters. *Glasg Nat* 2014; 26(1):103-104.
 36. Quignard JP, Tomasini JA. Mediterranean fish biodiversity. *Biol Mar Medit* 2000; 7(3):1-66.
 37. Roig A, Demestre M. Nota sobre la captura de dos *Zu cristatus* (Bonelli, 1820) en aguas del litoral catalán (Pisces, Trachipteridae). *Misc Zool* 1982; 6:152-154.
 38. Stipa MG, Longo F, Ammendolia G, Romeo T, Battaglia P. New data on *Trachipterus trachipterus* Gmelin, 1789 and *Zu cristatus* (Bonelli, 1820) (Pisces: Trachipteridae) from the Mediterranean Sea. *Acta Adriat* 2022; 63(1):65-74. doi: 10.32582/aa.63.1.7

39. Tiralongo F, Lillo AO, Tibullo D, Tondo E, Martire CL, D'Agnese R, Macali A, Mancini E, Giovos I, Coco S, Azzurro E. Monitoring uncommon and non-indigenous fishes in Italian waters: One year of results for the Alien Fish Project. *Reg Stud Mar Sci* 2019; 28:100606. doi: 10.1016/j.rsma.2019.100606
40. Tiralongo, F, Crocetta F, Riginella E, Lillo AO, Tondo E, Macali A, Mancini E, Russo F, Coco S, Paolillo G, Azzurro E. Snapshot of rare, exotic and overlooked fish species in the Italian seas: A citizen science survey. *J Sea Res* 2020; 164:101930. doi: 10.1016/j.seares.2020.101930
41. Zenetos A, Akel EHKh, Apostolidis C, Bilecenoglu M, Bitar G, Buchet V, Chalari N, Corsini-Foka M, Crocetta F, Dogrammatzi A, Drakulić M, Fanelli G, Giglio G, Imsiridou A, Kapisir K, Karachle PK, Kavadas S, Kondylatos G, Lefkaditou E, Lipej L, Mavrič B, Minos G, Moussa R, Prato E, Pancucci-Papadopoulou MA, Renda W, Ríos N, Rizkalla SI, Russo F, Servonnat M, Siapatis A, Sperone E, Theodorou JA, Tiralongo F, Tzovenis I. New Mediterranean Biodiversity Records (April 2015). *Mediterr Mar Sci* 2015; 16(1):266-284. doi: 10.12681/mms.1292